

## How big are atoms?

Atoms are very small. An atom and its parts are much smaller than a meter. The diameter of an atom is small  $10^{-10}$  (0.0000000001) meter, whereas an electron is smaller than  $10^{-18}$  (0.0000000000000001) meter. Comparatively, this means that an electron is 10 million times smaller than an atom! The diameter of a nucleon (a proton or neutron) is a distance that is equal to one fermi. This unit (equal to  $10^{-15}$  meter) is named for Enrico Fermi, an Italian-born physicist who studied the nucleus of the atom. For his work with neutrons, he received the Nobel Prize for physics in 1938.

**Most of the atom** You may be surprised to learn that most of the atom is actually empty space: If the atom was the size of your classroom, then the nucleus would be the size of a grain of sand in the center of the room.

## John Dalton and the atomic theory



As early as 400 BC, Greek philosophers proposed the atomic theory. This theory states that all matter is composed of tiny particles called atoms. Many centuries later, English chemist and physicist John Dalton (1766-1844) was one of the first scientists to set out to gather evidence for the idea. Dalton was a remarkable person. Born into a family too poor to send him to school, young John educated himself and, at age 12, became a schoolteacher. He grew to be one of the leading scientists of his time.

In 1808, Dalton published a detailed atomic theory that contained the following important points:

- 1 Each element is composed of extremely small particles called atoms.
  - 2 All atoms of a given element are identical.
  - 3 Atoms of different elements have different properties, including mass and chemical reactivity.
  - 4 Atoms are not changed by chemical reactions, but merely rearranged into different compounds.
  - 5 Compounds are formed when atoms of more than one element combine.
  - 6 A compound is defined by the number, type (element), and proportion of the constituent atoms.
- Dalton's atomic theory laid the groundwork for later atomic models, and over time, his original theory has been expanded and updated.

Particle	Diameter (meter)
atom	$10^{-10}$
nucleus	$10^{-14}$
proton	$10^{-15}$
neutron	$10^{-15}$
electron	$10^{-18}$

**Figure 18.3:** Diameter of atom and its subatomic particles

### Weather & atomic theory



One of John Dalton's records for 57 years, which led him to study how water into the air at an increased gas pressure these observations and from other he gathered evidence the structure of matter



## The changing model of the atom

The current model of the atom represents our current understanding of atomic structure. This model is one of a series of models constructed by people as they learned new information about atoms. New experimentation enabled people to update and change their ideas about how the atom is constructed.




The name *atom* comes from Democritus, a Greek philosopher (circa 460–370 BC) who proposed that matter is made up of small particles, which he called atoms, from the Greek word *atomos*, or indivisible. His model describes atoms as small particles that differ in size and shape, that combine in different configurations, and that are constantly in motion. Many of Democritus' ideas were based on logical thinking.

The idea that theories need to be supported by evidence—often gathered in carefully controlled experiments—became important in the 1600s. Then scientists began to design experiments to support and improve ideas proposed by earlier thinkers such as Democritus. John Dalton (see previous page) was a chemist who experimented with different gases. His careful measurements gave him repeatable evidence that matter is made up of atoms. His model of the atom is a tiny hard sphere (Figure 18.4).


The idea that atoms might contain smaller particles came about through a series of observations of cathode ray tubes, devices that were early versions of fluorescent and neon lights. Julius Plucker (1801–1844) and William Crookes, an English physicist and chemist (1832–1919), and his countryman and fellow physicist Joseph John Thomson (1856–1940) conducted many of these experiments. They proposed that different gases placed in the tubes generated streams of particles and conducted current.

In these experiments Thomson identified the electron, which carries a negative charge. Thomson knew that atoms were electrically neutral, so he proposed that the atom was a positive sphere with negative electrons embedded in it like raisins in a doughnut (Figure 18.5). The positive sphere and the negative electrons had an equal and opposite amount of charge, so the atom was neutral.

In England, physicists Ernest Rutherford (1871–1937), Hans Geiger (1882–1945), and Ernest Marsden (1889–1970), used high-speed, lightweight atoms called alpha particles (generated by radioactive material), to bombard very thin pieces of gold foil. Most of the particles passed through the foil and hit a screen behind it. But surprisingly, some of them bounced back. They must have hit areas of the foil with greater density!



**Figure 18.4:** Dalton's model of the atom. He thought atoms were tiny, hard spheres.



**Figure 18.5:** The Thomson model of the atom. The atom is a positive sphere with negative electrons embedded in it. Thomson discovered the electron.

Rutherford hypothesized that an atom must be made up of mostly empty space, allowing most of the alpha particles to pass through the foil. In the center of the atom, he suggested, was a tiny core called a nucleus, which contained positively-charged protons. This is where most of the mass must be found. The lighter electrons occupied the area between the nucleus and the edge of the atom. However, Rutherford did not have enough information to describe the electrons' location more fully.



Danish physicist Niels Bohr (1885-1962) used information about the nature of the emission of light by heated objects to update Rutherford's model. He described electrons as moving around the nucleus in fixed orbits that have a set amount of energy (figure 18.6). Bohr's model of the electron orbits is still used in many analyses of the atom. However, other 20th century experiments have shown that radiating waves can behave like particles in motion, and particles in motion can behave like waves.


In 1923, Louis de Broglie (1892-1987), a French physicist, showed how to analyze a moving particle as a wave. In 1926, Austrian physicist Erwin Schrödinger (1887-1961) built on de Broglie's work and treated electrons as three-dimensional waves. He developed a mathematical description of electrons in atoms that is called the quantum mechanical model of the atom. It is also called the electron cloud model, because his mathematical description cannot be described easily either in words or pictures, so a cloud represents the probability of electron position.

There still remained a serious problem with the atomic model, a problem Rutherford had identified so many years earlier: missing mass. In 1932, James Chadwick, an English physicist working in Rutherford's laboratory, finally solved the problem. He identified the third important subatomic particle, the neutron. Chadwick (1891-1974) based his work on earlier experiments by French physicists Irene and Frederic Joliot-Curie.

Understanding what is inside an atom has motivated many thousands of scientists and thinkers. What some of them discovered along the way changed the world, influencing not only theoretical spheres such as many of the sciences, philosophy, logic, and other areas, but also those subjects' practical applications. So many new technological developments of the late 20th century have been made possible by atomic research that the present era is often referred to as the "atomic age."

## 18.2 Compa


As you know, some s  
 elements. You alread  
 helium, silver, gold, a  
 exactly how does one  
 therapists, physicians, l  
 other medicines to be  
 how people figured ou  
 of the science. It br  
 chemists, who studied l



**Figure 18.6:** The Bohr model of the atom. Electrons move around the nucleus in fixed orbits.

## Atomic number

The number of Re  
 protons net  
 definition is an thi  
 element ato  
 can you change All  
 the number of diff  
 protons? prot  
 The  
 cher  
 The



the atomic  
 number?  
 atom:  
 Atom  
 11, 1  
 labora

**Figure 18.7:** The Schrödinger model of an atom is also called the electron cloud model. The cloud represents the probable locations of an electron.





### How big are atoms?

Atoms are very small. An atom and its parts are much smaller than a meter. The diameter of an atom is  $10^{-10}$  (0.0000000001) meter, whereas an electron is smaller than  $10^{-18}$  (0.000000000000000001) meter. Comparatively, this means that an electron is 10 million times smaller than an atom! The diameter of a nucleon (a proton or neutron) is a distance that is equal to one fermi. This unit (equal to  $10^{-15}$  meter) is named for Enrico Fermi, an Italian-born physicist who studied the nucleus of the atom. For his work with neutrons, he received the Nobel Prize for physics in 1938.

Most of the atom is empty space. You may be surprised to learn that most of the atom is actually empty space: If the atom was the size of your classroom, then the nucleus would be the size of a grain of sand in the center of the room.

### John Dalton and the atomic theory



As early as 400 BC, Greek philosophers proposed the atomic theory. This theory states that all matter is composed of tiny particles called atoms. Many centuries later, English chemist and physicist John Dalton (1766-1844) was one of the first scientists to set out to gather evidence for the idea. Dalton was a remarkable person. Born into a family too poor to send him to school, young John educated himself and, at age 12, became a schoolteacher. He grew to be one of the leading scientists of his time.

In 1808, Dalton published a detailed atomic theory that contained the following important points:

- 1 Each element is composed of extremely small particles called atoms.
  - 2 All atoms of a given element are identical.
  - 3 Atoms of different elements have different properties, including mass and chemical reactivity.
  - 4 Atoms are not changed by chemical reactions, but merely rearranged into different compounds.
  - 5 Compounds are formed when atoms of more than one element combine.
  - 6 A compound is defined by the number, type (element), and proportion of the constituent atoms.
- Dalton's atomic theory laid the groundwork for later atomic models, and over time, his original theory has been expanded and updated.

Particle	Diameter (meters)
atom	$10^{-10}$
nucleus	$10^{-14}$
proton	$10^{-15}$
neutron	$10^{-15}$
electron	$10^{-18}$

Figure 18.3: Diameters of atom and its subatomic particles.

### Weather & atomic theory




One of John Dalton's ideas was weather (he kept records for 57 years) and led him to study gas. He studied the evaporation of water into the air and was to understand that there increased gas pressure these observations of air and from other experiments he gathered evidence the structure of matter.



## The changing model of the atom

The current model of the atom represents our current understanding of atomic structure. This model is one of a series of models constructed by people as they learned new information about atoms. New information enabled people to update and change their ideas about how the atom is constructed.



The name *atom* comes from Democritus, a Greek philosopher (circa 460–370 BC) who proposed that matter is made up of small particles, which he called atoms, from the Greek word *atomos*, or indivisible. His model describes atoms as small particles that differ in size and shape, that combine in different configurations, and that are constantly in motion. Many of Democritus' ideas were based on logical thinking.

The idea that theories need to be supported by evidence—often gathered in carefully controlled experiments—became important in the 1600s. Then scientists began to design experiments to support or disprove ideas proposed by earlier thinkers such as Democritus. John Dalton (see previous page) was a chemist who experimented with different gases. His careful measurements gave him repeatable evidence that matter is made up of atoms. His model of the atom is a tiny hard sphere (figure 18.4).

The idea that atoms might contain smaller particles came about through a series of observations of cathode ray tubes, devices that were early versions of fluorescent and neon lights. Julius Plucker (1801–1888) and William Crookes, an English physicist and chemist (1832–1919), and his countryman and physicist Joseph John Thomson (1856–1940) conducted many of these experiments. They showed that different gases placed in the tubes generated streams of particles and conducted current.



In these experiments Thomson identified the electron, which carries a negative charge. Thomson knew that atoms were electrically neutral, so he proposed that the atom was a positive sphere with negative electrons embedded in it like raisins in a bun (figure 18.5). The positive sphere and the negative electrons had an equal and opposite amount of charge, so the atom was neutral.

In 1911 in England, physicists Ernest Rutherford (1871–1937), Hans Geiger (1882–1945), and Ernest Marsden (1889–1970), used high-speed, lightweight atoms called alpha particles (generated by radioactive material), to bombard very thin pieces of gold foil. Most of the alpha particles passed through the foil and hit a screen behind it. But surprisingly, some of them bounced back. They must have hit areas of the foil with greater density!

Diameter  
(meters)

$10^{-10}$

$10^{-14}$

$10^{-15}$

$10^{-15}$


$10^{-18}$

radii of  
atomic particles


atomic theory



Thomson's interest in these experiments led to the discovery of the electron. He proposed that atoms were electrically neutral, so he proposed that the atom was a positive sphere with negative electrons embedded in it like raisins in a bun (figure 18.5). The positive sphere and the negative electrons had an equal and opposite amount of charge, so the atom was neutral.



**Figure 18.4:** Dalton's model of the atom. He thought atoms were tiny, hard spheres.



**Figure 18.5:** The Thomson model of the atom. The atom is a positive sphere with negative electrons embedded in it. Thomson discovered the electron.



Rutherford hypothesized that an atom must be made up of mostly empty space, allowing most of the alpha particles to pass through the foil. In the center of the atom, he suggested, was a tiny core called a nucleus, which contained positively-charged protons. This is where most of the mass must be found. The lighter electrons occupied the area between the nucleus and the edge of the atom. However, Rutherford did not have enough information to describe the electrons' location more fully.




Danish physicist Niels Bohr (1885-1962) used information about the nature of the emission of light by heated objects to update Rutherford's model. He described electrons as moving around the nucleus in fixed orbits that have a set amount of energy (figure 18.6). Bohr's model of the electron orbits is still used in many analyses of the atom. However, other 20th century experiments have shown that radiating waves can behave like particles in motion, and particles in motion can behave like waves.

In 1923, Louis de Broglie (1892-1987), a French physicist, showed how to analyze a moving particle as a wave. In 1926, Austrian physicist Erwin Schrödinger (1887-1961) built on de Broglie's work and treated electrons as three-dimensional waves. He developed a mathematical description of electrons in atoms that is called the quantum mechanical model of the atom. It is also called the electron cloud model, because his mathematical description cannot be described easily either in words or pictures, so a cloud represents the probability of electron position.

There still remained a serious problem with the atomic model, a problem Rutherford had identified so many years earlier: missing mass. In 1932, James Chadwick, an English physicist working in Rutherford's laboratory, finally solved the problem. He identified the third important subatomic particle, the neutron. Chadwick (1891-1974) based his work on earlier experiments by French physicists Irene and Frederic Joliot-Curie.



Understanding what is inside an atom has motivated many thousands of scientists and thinkers. What some of them discovered along the way changed the world, influencing not only theoretical spheres such as many of the sciences, philosophy, logic, and other areas, but also those subjects' practical applications. So many new technological developments of the late 20th century have been made possible by atomic research that the present era is often referred to as the "atomic age."



**Figure 18.6:** The Bohr model of the atom. Electrons move around the nucleus in fixed orbits.

## 18.2 Compari

As you know, some scientists. You already know, silver, gold, aluminum, how does one determine, physicians, to better medicines to be

How people figured out things in science. It brings chemists, who studied h

### Atomic number

The number of protons determines an element's atomic number. All elements have a unique number of protons.

Can you change the number of protons? No, the number of protons in an atom is fixed.

What is the atomic number? The number of protons in an atom.

Atomic number is the number of protons in an atom. It is a fundamental property of an element.



**Figure 18.7:** Electrons in the Schrödinger model of the atom model is also called the electron cloud model. The cloud represents the probable locations of an electron.